## **IMPACT OF EVENT SCALE-REVISED**

## Daniel S. Weiss, PhD & Charles R. Marmar, MD

Instructions: Below is a list of difficulties people sometimes have after stressful life events.

Please read each item and then indicate how distressing each difficulty has been for you I PAST SEVEN DAYS with respect to				DURING THE how much		
Not at all=0, Little bit =1, Moderately=2, Quite a bit = 3, Extremely= 4						
Sr.No	Statement	0	1	2	3	4
1.	Any reminder brought back feelings about it					
2.	I had trouble staying asleep.					
3.	Other things kept making me think about it.					
4.	I felt irritable and angry.					
5.	I avoided letting myself get upset when I thought about it or was reminded of it.					
6.	I thought about it when I didn't mean to					
7.	I felt as if it hadn't happened or wasn't real					
8.	I stayed away from reminders about it.					
9.	Pictures about it popped into my mind.					
10	I was jumpy and easily startled.					
11	I tried not to think about it.					
12	I was aware that I still had a lot of feelings about it, but I didn't deal with					
	them.					
13	My feelings about it were kind of numb.					
14	I found myself acting or feeling like I was back at that time.					
15	I had trouble falling asleep.					
16	I had waves of strong feelings about it.					
17	I tried to remove it from my memory.					
18	I had trouble concentrating.					
19	Reminders of it caused me to have physical reactions, such as sweating,					
	trouble breathing.					
	I had dreams about it.					
	I felt watchful and on-guard.					
22	I tried not to talk about it.					

Avoidance Subscale = mean of items 5, 7, 8, 11, 12, 13, 17, 22

Intrusion Subscale = mean of items 1, 2, 3, 6, 9, 16, 20

Hyper arousal Subscale = mean of items 4, 10, 14, 15, 18, 19, 21

Note: The IES-R is not a diagnostic or screening tool for PTSD; rather, it relies on a patient's own report of symptoms and is used to gauge response no sooner than two weeks after a traumatic event, as well as to evaluate recovery.